

Scale setting in V+jets production

Christian Bauer

Lawrence Berkeley National Laboratory

Loopfest IX

Based on work in collaboration with Bjoern Lange

0905.4739

Motivation

- LO calculations are by now very easy
- Despite much progress on higher order calculations, many distributions still only available at LO
- Well known that distributions can have very different shape at LO and NLO
- For many processes, dominant effects at NLO arise from terms enhanced by logarithms of large ratios of scales

Motivation

- LO calculations are by now very easy
- Despite much progress on higher order calculations, many distributions still only available at LO
- Well known that distributions can have very different shape at LO and NLO
- For many processes, dominant effects at NLO arise from terms enhanced by logarithms of large ratios of scales

Can we resum these large logarithmic terms?

Why just scale setting?

- Large corrections arise from large logarithmic terms
- Can be resummed using IR evolution equations or effective field theory methods (SCET)
- Has allowed for much better predictions for many observables (thrust, Higgs production, Drell-Yan, ...)
- But most of these resummation have nothing to do with scale setting

Why just scale setting?

- Large corrections arise from large logarithmic terms
- Can be resummed using IR evolution equations or effective field theory methods (SCET)
- Has allowed for much better predictions for many observables (thrust, Higgs production, Drell-Yan, ...)
- But most of these resummation have nothing to do with scale setting

Can large logs be resummed by scale setting?

Why just scale setting?

- Large corrections arise from large logarithmic terms
- Can be resummed using IR evolution equations or effective field theory methods (SCET)
- Has allowed for much better predictions for many observables (thrust, Higgs production, Drell-Yan, ...)
- But most of these resummation have nothing to do with scale setting

Can large logs be resummed by scale setting?

If yes, what accuracy can be achieved?

Outline

- Quick overview of log resummation using SCET
- Case study: $pp \rightarrow Vj$
- Adding one extra jet: $pp \rightarrow Vjj$
- Discussion about adding additional jets

Log resummation using EFT's

Log resummation in SCET relies on RG evolution

Log resummation using EFT's

Log resummation in SCET relies on RG evolution

Derive RG Equation,
by taking $\mu \, d/d\mu$

$$\mu \frac{d}{d\mu} \frac{d\sigma_n^{\text{LL}}(\mu)}{d\Phi_n} = \gamma_n(\mu) \frac{d\sigma_n^{\text{LL}}(\mu)}{d\Phi_n}$$

Log resummation using EFT's

Log resummation in SCET relies on RG evolution

Derive RG Equation,
by taking $\mu \frac{d}{d\mu}$

Solve to find
(Δ_n = Sudakov factor)

$$\mu \frac{d}{d\mu} \frac{d\sigma_n^{\text{LL}}(\mu)}{d\Phi_n} = \gamma_n(\mu) \frac{d\sigma_n^{\text{LL}}(\mu)}{d\Phi_n}$$

$$\frac{d\sigma_n^{\text{LL}}(\mu)}{d\Phi_n} = \frac{d\sigma_n^{\text{LL}}(\mu_0)}{d\Phi_n} \Delta_n(\mu_0, \mu)$$

Log resummation using EFT's

Log resummation in SCET relies on RG evolution

Derive RG Equation,
by taking $\mu \frac{d}{d\mu}$

Solve to find
(Δ_n = Sudakov factor)

$$\mu \frac{d}{d\mu} \frac{d\sigma_n^{\text{LL}}(\mu)}{d\Phi_n} = \gamma_n(\mu) \frac{d\sigma_n^{\text{LL}}(\mu)}{d\Phi_n}$$

$$\frac{d\sigma_n^{\text{LL}}(\mu)}{d\Phi_n} = \frac{d\sigma_n^{\text{LL}}(\mu_0)}{d\Phi_n} \Delta_n(\mu_0, \mu)$$

Large logs: \longleftrightarrow Arise from IR divergences

RG evolution: \longleftrightarrow Uses UV divergences

Log resummation using EFT's

Log resummation in SCET relies on RG evolution

Derive RG Equation,
by taking $\mu \frac{d}{d\mu}$

Solve to find
(Δ_n = Sudakov factor)

$$\mu \frac{d}{d\mu} \frac{d\sigma_n^{\text{LL}}(\mu)}{d\Phi_n} = \gamma_n(\mu) \frac{d\sigma_n^{\text{LL}}(\mu)}{d\Phi_n}$$

$$\frac{d\sigma_n^{\text{LL}}(\mu)}{d\Phi_n} = \frac{d\sigma_n^{\text{LL}}(\mu_0)}{d\Phi_n} \Delta_n(\mu_0, \mu)$$


Large logs: \longleftrightarrow Arise from IR divergences

RG evolution: \longleftrightarrow Uses UV divergences


What do large logs have to do with UV of theory?

Log resummation using EFT's

Logarithms related to IR divergences in theory




Divergences from loop integrations
 $1/\epsilon + \dots$




Divergences from phase space integrations
 $\int_0^1 dk_g (k_g)^{-1-\epsilon} = -1/\epsilon + \dots$

Log resummation using EFT's

Logarithms related to IR divergences in theory



Divergences from loop integrations
 $1/\epsilon + \dots$



Divergences from phase space integrations
 $\int_0^1 dk_g (k_g)^{-1-\epsilon} = -1/\epsilon + \dots$

Restrictions on phase space give rise to logarithmic remainders
 $0 < k_g < \mu \Rightarrow -1/\epsilon + \log(\mu)$

$$\sigma_V + \sigma_R = \log(\mu) + \dots$$

Log resummation using EFT's

Effective theories are defined to reproduce the IR physics of underlying theory

Log resummation using EFT's

Effective theories are defined to reproduce the IR physics of underlying theory

If EFT contains no intrinsic mass scales, it only depends on Λ_{IR} and Λ_{UV}

Log resummation using EFT's

Effective theories are defined to reproduce the IR physics of underlying theory

If EFT contains no intrinsic mass scales, it only depends on Λ_{IR} and Λ_{UV}

In pure dim-reg $\Lambda_{\text{IR}} \rightarrow 0$ and $\Lambda_{\text{UV}} \rightarrow \infty$

Log resummation using EFT's

Effective theories are defined to reproduce the IR physics of underlying theory

If EFT contains no intrinsic mass scales, it only depends on Λ_{IR} and Λ_{UV}

In pure dim-reg $\Lambda_{\text{IR}} \rightarrow 0$ and $\Lambda_{\text{UV}} \rightarrow \infty$

No scale in problem \Rightarrow result in EFT is 0

Log resummation using EFT's

Effective theories are defined to reproduce the IR physics of underlying theory

If EFT contains no intrinsic mass scales, it only depends on Λ_{IR} and Λ_{UV}

In pure dim-reg $\Lambda_{\text{IR}} \rightarrow 0$ and $\Lambda_{\text{UV}} \rightarrow \infty$

No scale in problem \Rightarrow result in EFT is 0

$$\Rightarrow \text{Log}(\Lambda_{\text{IR}}) = - \text{Log}(\Lambda_{\text{UV}})$$

Log resummation using EFT's

Effective theories are defined to reproduce the IR physics of underlying theory

If EFT contains no intrinsic mass scales, it only depends on Λ_{IR} and Λ_{UV}

In pure dim-reg $\Lambda_{\text{IR}} \rightarrow 0$ and $\Lambda_{\text{UV}} \rightarrow \infty$

No scale in problem \Rightarrow result in EFT is 0

$$\Rightarrow \text{Log}(\Lambda_{\text{IR}}) = - \text{Log}(\Lambda_{\text{UV}})$$

IR dependence of full theory can be extracted from UV dependence of EFT

Log resummation using EFT's


Log resummation using EFT's

- Consider problem with many widely separated scales
 $Q \gg \mu_1 \gg \dots \gg \mu_n \gg \Lambda$

Q
 μ_1
 \dots
 μ_n
 Λ


Log resummation using EFT's

- Consider problem with many widely separated scales
 $Q \gg \mu_1 \gg \dots \gg \mu_n \gg \Lambda$
- At Q match QCD to SCET setting other scales to zero




Log resummation using EFT's

- Consider problem with many widely separated scales
 $Q \gg \mu_1 \gg \dots \gg \mu_n \gg \Lambda$
- At Q match QCD to SCET setting other scales to zero
- Evolve from Q to μ_1 using RGE




Log resummation using EFT's

- Consider problem with many widely separated scales
 $Q \gg \mu_1 \gg \dots \gg \mu_n \gg \Lambda$
- At Q match QCD to SCET setting other scales to zero
- Evolve from Q to μ_1 using RGE
- Match onto new theory obtained by setting $\mu_1 \rightarrow \infty$




Log resummation using EFT's

- Consider problem with many widely separated scales
 $Q \gg \mu_1 \gg \dots \gg \mu_n \gg \Lambda$
- At Q match QCD to SCET setting other scales to zero
- Evolve from Q to μ_1 using RGE
- Match onto new theory obtained by setting $\mu_1 \rightarrow \infty$
- Repeat to remove all scales




Case Study: $pp \rightarrow V+j$

Study the p_T distribution at large p_T



Case Study: $pp \rightarrow V+j$

Study the p_T distribution at large p_T




Can these diverging curves be reconciled using resummation?

Case Study: $pp \rightarrow V+j$

Study the p_T distribution of jet, in region $p_T \gg m_W$

Three scales in problem: $p_T \gg m_W \gg \Lambda$

Integrate them out one by one




Case Study: $pp \rightarrow V+j$

Study the p_T distribution of jet, in region $p_T \gg m_W$

Three scales in problem: $p_T \gg m_W \gg \Lambda$

Integrate them out one by one




Resums all logs of form m_W^2/p_T^2

Case Study: $pp \rightarrow V+j$

Study the p_T distribution of jet, in region $p_T \gg m_W$

Three scales in problem: $p_T \gg m_W \gg \Lambda$


Integrate them out one by one




Resums all logs of form m_W^2/p_T^2

But what is really happening at scale m_V ?

Case Study: $pp \rightarrow V+j$




Case Study: $pp \rightarrow V+j$




Case Study: $pp \rightarrow V + j$

Integrate
out off-
shell quark



$\infty, 0, 0$

m_V




$\infty, \infty, 0$


Make V
boson static

Case Study: $pp \rightarrow V+j$

Integrate
out off-
shell quark



m_V




Make V
boson static


Two different operators, but both have same strongly interacting field content

Case Study: $pp \rightarrow V + j$

Integrate
out off-
shell quark



m_V




Make V
boson static

Two different operators, but both have same strongly interacting field content

Implies that the leading UV divergences are same for both operators

LL running above m_V and below m_V the same

Case Study: $pp \rightarrow V + j$



Two different operators, but both have same strongly interacting field content

Implies that the leading UV divergences are same for both operators

LL running above m_V and below m_V the same

No leading log dependence on scale m_V

Case Study: $pp \rightarrow V+j$

Only relevant scales at LL are Q and Λ

Running in EFT is given by AP evolution kernels

All leading logs resummed by choosing $\mu_F = p_T$

Case Study: $pp \rightarrow V+j$

Only relevant scales at LL are Q and Λ

Running in EFT is given by AP evolution kernels

All leading logs resummed by choosing $\mu_F = p_T$

Why not true beyond LL?


Case Study: $pp \rightarrow V+j$

Only relevant scales at LL are Q and Λ

Running in EFT is given by AP evolution kernels

All leading logs resummed by choosing $\mu_F = p_T$

Why not true beyond LL?



Different energy, will affect
subleading divergences

Results only
correct at LL
accuracy, does
not hold beyond

Case Study: $pp \rightarrow V+j$

- This is of course well known
- Most NLO calculations use a “dynamical scale”
 $\mu^2 = p_T^2 + m_W^2$
- Important result is
 - This can be shown to be correct at LL accuracy
 - Can be shown through a simple EFT analysis


Case Study: $pp \rightarrow V+j$

- This is of course well known
- Most NLO calculations use a “dynamical scale”
 $\mu^2 = p_T^2 + m_W^2$
- Important result is
 - This can be shown to be correct at LL accuracy
 - Can be shown through a simple EFT analysis

Can extend this result to higher number of jets and eventually to very different processes

One extra jet: $pp \rightarrow V + jj$

Consider $Q \sim p_T \gg m_W \gg \Lambda$




As before, nothing happens for LL at m_V

Both operators have same strong fields

Should again choose $\mu_F = p_T$

One extra jet: $pp \rightarrow V + jj$


One side note:



This matching assumes that $p_2^2 \gg p_1^2$

One extra jet: $pp \rightarrow V + jj$

One side note:




This matching assumes that $p_2^2 \gg p_1^2$

This implies that the
to jets should be back-
to back in Φ


One extra jet: $pp \rightarrow V + jj$

One side note:




This matching assumes that $p_2^2 \gg p_1^2$

This implies that the
to jets should be back-
to back in ϕ




One extra jet: $pp \rightarrow V + jj$

Compare again to fixed order calculations



One extra jet: $pp \rightarrow V + jj$


Compare again to fixed order calculations



Again, with new scale choice, much better agreement between LO and NLO calculations


Adding extra jet: $pp \rightarrow V + jjj$

- Nothing in this discussion knew anything about the number of final state jets
- In general, should choose scale $\mu = Q_{\text{QCD}}$, where Q_{QCD} denotes scale at which jets are produced



Adding extra jet: $pp \rightarrow V + jjj$

- Nothing in this discussion knew anything about the number of final state jets
- In general, should choose scale $\mu = Q_{\text{QCD}}$, where Q_{QCD} denotes scale at which jets are produced



At time of our paper, no NLO calculation available


Precise scale choice

NLO calculations now exist!

Blackhat, 0907.1984

Clearly, there
are various
ways to
define Q_{QCD}

Can not use
EFT's right now
to prefer one
over another



Conclusions

- Large logarithms often plague fixed order calculations
- In some cases, large logarithms arise only from unfortunate scale choices
- EFT's allow to understand logarithms and therefore appropriate scale choices naturally
- Have given examples in $pp \rightarrow V + \text{jets}$ where proper scale choice can significantly improve convergence of $LO \rightarrow NLO$
- Works very well, but precise scale choice can not be predicted
- Expect that same ideas hold for many other processes